Food and Drug Administration, HHS

(parasitic stages) (Hypoderma bovis, H. lineatum); lice (Linognathus vituli, Haematopinus eurysternus, Solenopotes capillatus); mites (Psoroptes ovis (syn. P. communis var. bovis), Sarcoptes scabiei var. bovis); and for control of infections of D. viviparus and O. radiatum for 28 days after treatment; O. ostertagi, T. axei, and C. punctata for 21 days after treatment; and H. placei and C. oncophora for 14 days after treatment.

(3) Limitations. For subcutaneous use Not for only. intravenous intramuscular use. Do not treat cattle within 49 days of slaughter. Because a withdrawal time in milk has not been established, do not use in female dairy cattle of breeding age. Do not use in other animal species because severe adverse reactions, including fatalities in dogs, may result. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

[55 FR 38984, Sept. 24, 1990, as amended at 62 FR 14302, Mar. 26, 1997; 62 FR 63271, Nov. 28, 1997; 64 FR 26671, May 17, 1999; 69 FR 31735, June 7, 2004; 72 FR 27734, May 17, 2007; 77 FR 64717, Oct. 23, 2012]

§ 522.1204 Kanamycin.

- (a) Specifications. Each milliliter of solution contains 50 or 200 milligrams (mg) of kanamycin as kanamycin sulfate
- (b) Sponsor. See No. 054771 in § 510.600(c) of this chapter.
- (c) Conditions of use in dogs and cats—(1) Amount. Administer by subcutaneous or intramuscular injection 5 mg per pound of body weight per day in equally divided doses at 12-hour intervals.
- (2) Indications for use. For the treatment of bacterial infections due to kanamycin sensitive organisms in dogs and cats.
- (3) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[79 FR 16190, Mar. 25, 2014]

§ 522.1222 Ketamine.

(a) Specifications. Each milliliter contains ketamine hydrochloride equivalent to 100 milligrams (mg) ketamine base activity.

- (b) Sponsors. See Nos. 000859, 026637, 054628, 054771, 061690, and 063286 in $\S 510.600(c)$ of this chapter.
- (c) Special considerations. Federal law restricts this drug to use by or on the order of a licensed veterinarian.
- (d) Conditions of use—(1) Cats—(i) Amount. 5 to 15 mg/pound body weight intramuscularly, depending on the effect desired.
- (ii) *Indications for use*. For restraint or as the sole anesthetic agent in diagnostic or minor, brief surgical procedures that do not require skeletal muscle relaxation.
- (2) Subhuman primates—(i) Amount. 3 to 15 mg/kilogram body weight intramuscularly, depending upon the species, general condition, and age of the subject.
 - (ii) *Indications for use*. For restraint.

 $[67\ FR\ 17283,\ Apr.\ 10,\ 2002,\ as\ amended\ at\ 73\ FR\ 8192,\ Feb.\ 13,\ 2008;\ 74\ FR\ 36111,\ July\ 22,\ 2009;\ 74\ FR\ 66573,\ Dec.\ 16,\ 2009;\ 75\ FR\ 10167,\ Mar.\ 5,\ 2010;\ 78\ FR\ 21060,\ Apr.\ 9,\ 2013.\ Redesignated\ and\ amended\ at\ 79\ FR\ 16191]$

§ 522.1223 Ketamine, promazine, and aminopentamide.

- (a) Specifications. Each milliliter of solution contains ketamine hydrochloride equivalent to 100 milligrams (mg) ketamine base activity, 7.5 (mg) of promazine hydrochloride, and 0.0625 mg of aminopentamide hydrogen sulfate.
- (b) Sponsor. See No. 054771 in $\S510.600$ (c) of this chapter.
- (c) Conditions of use in cats—(1) Amount. Administer by intramuscular injection 15 to 20 mg ketamine base per pound of body weight, depending on the effect desired.
- (2) Indications for use. It is used in cats as the sole anesthetic agent for ovariohysterectomy and general surgery.
- (3) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[79 FR 16191, Mar. 25, 2014]

§ 522.1225 Ketoprofen.

- (a) *Specifications*. Each milliliter of solution contains 100 milligrams (mg) of ketoprofen.
- (b) *Sponsor*. See No. 054771 in §510.600(c) of this chapter.